

**Dr. Trevor M. Brooks MD, FRCSC
Plastic, Hand & Cosmetic Surgery**

2- 1335 Trans Canada Way SE, Medicine Hat, AB, T1B 1J1
Tel: 403-487-0460 Fax: 403-487-0462
www.drtrevorbrooks.ca
E-mail: info@drtrevorbrooks.ca

Call our office or go to a Walk in Clinic if you have:

- Fever over 37 degrees
- Yellow, green or foul-smelling drainage
- A large red area around the incision
- An allergic reaction to the medications, or dressings (this could be shortness of breath, a rash/redness, hives, etc.)

Abdominoplasty Surgery Post Operative Care Instructions

Dressing/Bandages: Your dressings applied in the operating room should remain on, dry and in tact for 5-7 days following surgery. You should not shower during this time.

Your binder should be worn at all times for 6 weeks. If you take it off to wash it, please REST while you are not wearing it.

You will see Dr. Brooks at his office 7-8 days after your surgery for your first dressing change. You may shower the day after he sees you. Dry the area by patting with a clean towel or allowing to air dry. After each shower place a very thin layer of Polysporin on the incisions. If you have Steri-strips, let them fall off on their own.

There will be bloody drainage on the dressings – this is normal.

If bleeding occurs, hold constant & firm pressure for 10 mins with a clean towel. If bleeding continues and cannot be stopped, seek emergency care and then contact Dr. Brooks to inform him.

Do not submerge the area in water for 3 weeks (no baths, hot-tubs, pools, ocean/lake water). Your sutures are dissolvable and submerging them in water could open your incision up!

You will see the nurses at Day Medicine (at MHRH 2nd floor) for your drain removals once drainage is Less than < 20 ml over 24 hours. You will be taught how to monitor your drain output before you are discharged from the hospital on the day of your surgery.

Medication: Take your regular medications as prescribed (unless Dr. Brooks has specifically instructed you otherwise). Typically a pain medication, an antibiotic and a muscle relaxant will be prescribed after surgery.

DO NOT DRIVE WHILE TAKING THE PAIN MEDICATION.

Make sure to finish the entire Antibiotic prescription.

Once done the pain prescription, you may use Acetaminophen (TYLENOL) unless Dr. Brooks has instructed you otherwise.

AVOID ADVIL (ibuprofen) and inflammatory meds (Alleve) for the first 4 days.

Return to Work: Most patients are able to return to work 3-4 weeks after the surgery. People with jobs that require heavy lifting may require more time before returning to full duties. We can provide you with a work note.

Your Follow Up Appointment: 7 days after your surgery Dr. Brooks will want to see you again. To ensure you get in within this time frame please call promptly after your surgery to set up the appointment 403-487-0460

Activity: You should take it easy for the first week, but you NEED to be UP and walking around 4-5 times a day to decrease the risk of blood clots.

Swelling is normal – for the 1st week keep your torso elevated with pillows and your hips at an angle like in a lazy-boy chair while sleeping on either your back or side. Or feel free to actually sleep in a lazy-boy reclining chair. Continue on your back or side for the first 4 weeks. If comfortable, you may sleep on your stomach after 4 weeks.

Beginning the second week you may begin to be fully upright as your pain tolerance permits.

Avoid lifting, pushing, pulling any object heavier than 10 lbs for the first 4 weeks (for reference a 4L milk jug is 9 lbs).

Avoid sexual activity for the first 2 weeks, and then be careful for the next 2 weeks.

Do not do sports, heavy house/yard work, or use exercise equipment until 4-6 weeks after surgery, or Dr. Brooks has given you the go ahead.

You may begin scar massage at 3 weeks (if no open areas to the incision). Use firm pressure & cream (Vit A, Vit E or Aloe are all excellent choices), and push against the direction of the scar (perpendicular).

A high quality Vitamin & silicone scar gel is available at Dr. Brooks's office, but there are many available at most pharmacies over the counter.

Diet: Resume your regular diet. Drink plenty of fluids, but stay away from alcoholic drinks for 1 week and limit caffeinated drinks for 1 week.

Avoid Smoking for 3 months before and 6 weeks after surgery for better wound healing. Nicotine constricts blood vessels.

Sunshine: After 3 weeks if your surgical site is to be exposed to sun, apply sunscreen SPF 45 or higher for at least 1 year to prevent the scar from changing to a darker colour.